

chapter 2

- 4) The unitarity triangle is not unique to the B^0 meson, there are in fact six of them. Construct the unitarity triangle for the K^0 system, i.e. the triangle that follows from the orthogonality condition on the 2nd row and 1st column, from $V^\dagger V = \mathbb{1}$, see Eq. (2.1).
- Write out the orthogonality condition of the 2nd row and 1st column, from $V^\dagger V = \mathbb{1}$ (ignore $\mathcal{O}(\lambda^7)$ terms).
 - Represent the condition as a triangle in the complex plane.
 - Rescale and rotate the triangle such that the basis of the triangle is formed by the points (0,0) and (1,0).
 - What are the coordinates of the apex of the triangle in terms of the Wolfenstein parameters?
 - Using the values of A, λ, ρ and η ($A = 0.82, \lambda = 0.22, \rho = 0.25$ and $\eta = 0.35$), make a drawing of the K^0 triangle that accurately reflects the shape of the triangle (i.e. the height-to-width ratio).
 - Calculate the surface area of the K^0 triangle in terms of the Wolfenstein parameters and compare it to the surface area of the B^0 triangle.
 - If we now include $\mathcal{O}(\lambda^6)$ -terms in the expression of the CKM-matrix with phases, on which element is an extra phase missing? (See Eq. 2.16):

$$V_{CKM, \text{Wolfenstein}} = \begin{pmatrix} |V_{ud}| & |V_{us}| & |V_{ub}|e^{-i\gamma} \\ -|V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}|e^{-i\beta} & -|V_{ts}|e^{i\beta_s} & |V_{tb}| \end{pmatrix} + \mathcal{O}(\lambda^5)$$

- 5) The uncertainties on the measurements of moduli of the CKM-elements, $|V_{ij}|$ is larger than in expression Eq. (2.6):

$$V_{CKM} = \begin{pmatrix} 0.97419 & 0.2257 & 0.00359 \\ 0.2256 & 0.97334 & 0.0415 \\ 0.00874 & 0.0407 & 0.999133 \end{pmatrix} \pm \begin{pmatrix} 0.00022 & 0.0010 & 0.00016 \\ 0.0010 & 0.00023 & 0.0011 \\ 0.00037 & 0.0010 & 0.000044 \end{pmatrix} \quad (6.2)$$

- Explain why a 4th generation can affect the result.
- What would be your guess of a CKM-matrix with 4 generations (in terms of λ only) ?
- Check if your guess is compatible with observations, by calculating one unitarity relation numerically, taking into account the uncertainty on the measured moduli $|V_{ij}|$.