

ECFA/DESY AMSTERDAM

e^-e^- parallel session:

THE NEED FOR COMPLEMENTARITY OF e^+e^-
AND e^-e^- COLLISIONS:

Recall that the $|in\rangle$ states of best definition

are $|e_L e_L\rangle$, $|e_L e_R\rangle$, $|e_R e_R\rangle$.


There is no other state so fully parameterized.

GIVEN THAT THE $|e^+e^-\rangle$ COLLIDER IS MOSTLY
VAUNTED FOR ITS "VACUUM" QUANTUM
NUMBERS, THERE IS A STRONG CASE TO
BE MADE FOR PROVIDING BOTH AT THE FIRST
STAGE OF LINEAR COLLIDER OPERATION.

RECALL THAT THE 1995 INTERNATIONAL
LINEAR COLLIDER TECHNICAL REVIEW COMMITTEE
REPORT STATED:

"Since the e^-e^- collider requires only minor changes to the hardware of the e^+e^- machine and detector, its programme could be pursued during the first phase of the facility...."

- International Linear Collider Technical Review Committee Report - 1995



THE ONLY REALLY PURE FIELD AT
OUR DISPOSAL IS

e_L, e_R

NOT AVAILABLE IN e^+
 γ

hadron beams

(also, to a certain extent, μ 's; but we
cannot switch their polarization).

$|\vec{P}_e| = 80\%$ is routinely reached
in SLQ running at SLAC

- There is no known cath' limit'n
- Theoretical estimates put
attainable $|\vec{P}|_e$ values at $\leq 95\%$

$|\vec{P}|_{\mu}$ is a much harder task

NO PROJECT BEYOND 65%
IS BEING ENVISIONED

... UNIQUE QUANTUM NUMBERS FOR e^-e^- INITIAL STATE.

A POTENT TOOL FOR

CONSERVATION STUDIES IN
SEARCH OF

⇒ FUNDAMENTAL SYMMETRIES

⇒ EXTENDED GROUP STRUCTURES
BEYOND THE STANDARD MODEL

TO WIT: e^-e^- INITIAL STATE HAS

$Q_u = -2$	
$L = +2$	$L_e = +2$
$I_w = -1$	
$Y_w = -2$	
$S_2 = 1, 0, -1$ (easily tunable)	

THESE PROPERTIES PERMIT STUDIES OF

- THE MEANING OF LEPTON #

$$\left(\sum (B-L) = \text{const?} \right)$$

- m_{Dirac} vs. m_{Majorana} FOR ν_e

A CLOSER LOOK AT THE DESIGN FEATURES
SHOWS THAT

THE JLC, TESLA, NLC, CLIC
ARE ALL NATURAL e^-e^-
COLLIDERS

IF

WE PROVIDE THE NEEDED
COMPLEMENTARY FEATURES
FROM THE START!

THE SAME HOLDS FOR I.R., DETECTOR.

THE ACCESSIBILITY OF

①

- HIGH DEGREES OF POLARIZATION
- EASY HELICITY REVERSAL

FOR BOTH INCOMING BEAMS

IS ESSENTIAL FOR ENHANCEMENT OF SIGNAL
AND SUPPRESSION OF BACKGROUND

→ EASY CHOICE OF CHIRAL COUPLINGS

②

- DISTINCTIVE CHOICE OF
INCOMING QUANTUM #^{'s}
FOR INITIAL STATE:

	Q_{ee}	S_2	L	L_2	I_3^W	Y^W
$e_L^- e_L^-$	-2	+1	2	2	-1	-2
$e_L^- e_R^-$	-2	0	2	2	$-\frac{1}{2}$	-3
$e_R^- e_R^-$	-2	-1	2	2	0	-4

PERMITS PROBING BEYOND THE CAPABILITIES
OF LHC AND e^+e^- COLLIDERS

PHYSICS PROMISE

Same-sign colliders are
COMPLEMENTARY
to Opposite-sign colliders.

→ MØLLER SCATTERING **MARCIANO**
2 ANOMALIES
(→ Z' , $\sin^2 \Theta_W$)

HIGGS (minimal, MSSM)

EXTENDED HIGGS SECTOR **J. Gunion**

→ SUSY **J. Feng, M. Peskin, T. Han**

STRONG EWSB

→ COMPOSITENESS **T. Barklow**

HEAVY MAJORANA NEUTRINOS **H.-M. Min**

DILEPTON GAUGE BOSONS **FRAMPTON**

LEPTOQUARKS **T. Rizzo**

ANOMALOUS GAUGE COUPLINGS **CUYPERS**

e^+e^- PROVIDES A LABORATORY
FOR PROBING

- THE S.M. TO ITS LIMITS
- HIGHER SYMMETRIES BEYOND
THE REACH OF e^+e^- , pp

THE ACCESSIBILITY OF SUCH CLEAN,

- "EXOTIC" QUANTUM NUMBERS

- A CLEAR CHOICE OF CHIRAL
COUPLINGS

THROUGH INSTANTLY SWITCHABLE

$e_L e_L$, $e_L e_R$, $e_R e_R$

COMPOSITIONS

MAKES e^+e^- A UNIQUE TOOL FOR PROBING

DETAILED FEATURES THAT ARE

} HARDER TO DETECT IN e^+e^- , pp
COMPLEMENTARY TO WHAT CAN
BE DISCOVERED OR EXPLORED
IN OTHER CHANNELS

EXAMPLES: ANOMALOUS GAUGE COUPLINGS
STRONG WW SECTOR
EXTENDED HIGGS MODELS

WHAT'S SPECIAL ABOUT e^-e^- VS. e^+e^- ?

- BEAM CURRENT MAY BE HIGHER
- "ANTI-PINCH" - BEAM-BEAM DISRUPTION

→ MAY LEAD TO CONSIDERATION OF
INSERTION OF A PLASMA LENSE

(BEAM-GAS BACKGROUNDS?)

- e^-e^- IS THE NECESSARY STARTING POINT
FOR e^-x , xx PROGRAM

→ THIS UNDERLINES THE NEED TO ESTABLISH e^-e^- FROM THE START,

BUT, DETECTOR ISSUES HAVE MORE IN
COMMON WITH e^+e^- THAN WITH e^-x , xx

- CROSSING ANGLE?

e^- WILL NOT BE GUIDED INTO

e^- BEAM OPTICS ON OTHER SIDE
OF I.R. (LIKE e^+e^- !)

- BEAM DISPOSITION, BEAM DUMPS ARE
TO BE CONFIGURED SPECIALLY!

- PERMANENT-MAGNET SCENARIO: ROTATION

e^- POLARIZATION

A VITAL INGREDIENT IN ESSENTIALLY ALL
THE MAIN PHYSICS MOTIVATIONS WE DISCUSSED

MAJOR RECENT DEVELOPMENTS

(HAVE PUT S.L.C. BACK INTO THE
DRIVER'S SEAT...)

WE EXPECT TO POLARIZE BOTH INCOMING
 e^- BEAMS "AT WILL" WITH $|\vec{P}| \geq 80\%$ ^{prob'ly 90%}
(superior to μ beams that hail from π, K decays!)

QUESTION: CAN THIS POLARIZATION BE
MAINTAINED FOR BOTH BEAMS IN $10^{19} e^- \text{ cm}^{-3}$
BUNCH-BUNCH INTERACTIONS?

DEPOLARIZATION SIMULATIONS
INDICATE ONLY SMALL EFFECTS
→ MORE WORK NEEDED.

"EXOTIC" QUANTUM #'s

easy high polarizability and helicity reversal
permit definition of

	Q_e	S_e	L	L_e	I_3^W	Y^W
$\bar{e}_L e_L$	-2	1	2	2	-1	-2
$\bar{e}_L e_R$	-2	0	2	2	$-\frac{1}{2}$	-3
$\bar{e}_R e_R$	-2	-1	2	2	0	-4

- signal identification, enhancement
- background reduction
- signal interpretation

for compositeness scales well into the 100 TeV e^-e^- colliders.





Figure 8: a) The sensitivity of Møller vs. Bhabha scattering below and above the Z resonance, in the absence of polarization of the incoming beams, to a compositeness scale Λ [TeV]. b) The corresponding sensitivity comparison in the case where the incoming electron (but not positron) beams are 90% polarized. The luminosity is chosen such that the statistics at each energy corresponds to the integrated values reached at PETRA for the compositeness limits set there.

• COMPOSITENESS OF ELECTRONS

TIM BARKLOW

AT SOME NEW-INTERACTION ENERGY SCALE Λ ,
ELECTRON COMPOSITENESS MAY MANIFEST ITSELF.

STANDARD LAGRANGIAN WITH THIS NEW FIELD:



$$\mathcal{L}_{\psi\psi} = \frac{g^2}{2\Lambda^2} \left\{ \eta_{LL} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L + \eta_{RR} \dots \right. \\ \left. + \eta_{RL} \bar{\psi}_R \gamma_\mu \psi_R \bar{\psi}_L \gamma^\mu \psi_L \right\}$$

THE EXPLICIT TREATMENT SHOWS THAT THE
NEW CONTACT INTERACTION IS CONSIDERABLY MORE
EFFECTIVE FOR e^-e^- THAN FOR e^+e^-

PARTICULARLY WHEN THE e^- ARE STRONGLY
POLARIZED


Λ - NEW-INTERACTION SCALES ACCESSIBLE
ARE MUCH HIGHER



HIGGS BOSON PRODUCTION:

STANDARD HIGGS SCALAR

FROM ZZ FUSION



EXCELLENT CHANNEL FOR DISCOVERY,
DETAILED STUDY FOR $\sqrt{s} \gtrsim 2.5 m_H$


$\Rightarrow \sigma_{\text{tot}}$ SATURATES FOR ENERGIES BEYOND
THRESHOLD

$\Rightarrow e^-e^-$ FINAL STATE CAN OFTEN BE DE-
TECTED (UNLIKE $e^+e^- \rightarrow \nu\bar{\nu}H$)

AND INVISIBLE HIGGS DECAYS

FROM "MISSING CENTRAL MASS"

$\Rightarrow$ PRODUCTION CROSS-SECTION HAS LITTLE
DEPENDENCE ON e^- POLARIZATION,
BUT



REFERENCES:


- K. HIKASA (1988)
- T. HAN (1995)
- P. MINIKOWSKI (1996)

Higgs boson production at e^+e^- colliders :


$\left\{ \begin{array}{l} \text{minimal } H^0 \\ \text{non } h^0 \end{array} \right\}$

1. Single Higgs production:

via $\gamma\gamma$ and ZZ fusion




basically a
"missing central
mass"
measurement





$\left\{ \begin{array}{l} \text{add} \\ \text{CP-odd} \\ A^0 \end{array} \right\}$

vs. e^+e^- collider standard graph.



$$\sigma \sim \frac{1}{s}$$

H. Haber
(1997)



a) Central production of neutral bosons by gamma-gamma or ZZ fusion. b) same as a), but the lepton loop also permits the production of CP-odd bosons. c) Production cross section for the kinematical parameters given: the saturation at higher energies is obvious; a reasonable detector-originated cut of 5 degrees is entirely acceptable. d) Simulated signal (on a log scale). for the parameters given, irrespective of the boson decay modes. It can almost act as a "Higgs trigger".

P. Minkowski, 20 97

$$e^- e^- \rightarrow (e^- e^-) H$$



19

DETECTOR

CUT

AT

$\Theta = \begin{cases} 5^\circ \\ 10^\circ \end{cases}$



SPECTACULAR ADVANTAGES OF
 e^-e^- COLLISIONS FOR SUSY STUDIES:

→ $\tilde{e}$ ARE SCALARS!

→ $m(\tilde{e}_R) \neq m(\tilde{e}_L)$!

SEE SUSY PARALLEL SESSION
TOMORROW !