

## chapter 4 (lecture 4)

- 7) Most recent measurements on CP violation are performed in the  $B^0$  system at the B-factories in California (*BaBar* experiment) and in Japan (*Belle* experiment). An interesting category are  $B^0$ -decays to CP-eigenstates,  $f = \bar{f}$ , because both the  $B^0$  and the CP-conjugated  $B$ , the  $\bar{B}^0$  can decay to this same final state.
- (a) What do the assumptions  $|q/p| = 1$  and  $|A_f| = |\bar{A}_f|$  imply?
- (b) If we assume  $|q/p| = 1$  and  $|A_f| = |\bar{A}_f|$ , the expression for  $A_{CP}$ ,

$$A_{CP}(t) = \frac{\Gamma_{P^0(t) \rightarrow f} - \Gamma_{\bar{P}^0(t) \rightarrow f}}{\Gamma_{P^0(t) \rightarrow f} + \Gamma_{\bar{P}^0(t) \rightarrow f}},$$

simplifies considerably. Write the expression for  $A_{CP}(t)$  if we in addition assume  $\Delta\Gamma \sim 0$ .

- (c) Under these three assumptions, is there CP-asymmetry at each value of  $t$ ?
- (d) Under these three assumptions, what is the time-integrated CP-asymmetry?
- (e) Which of these three assumptions is valid for each final state of the  $B^0$ ?
- (f) Write the general expression for  $A_{CP}(t)$  if we assume  $B^0$ -decays only.
- 8) To have observable CP violation in a process resulting from two interfering amplitudes one must have a phase difference between the amplitudes that changes under CP conjugation. The actual requirement is slightly more specific. The goal of this exercise is to formulate the more exact requirement.
- (a) Given a decay process that can proceed through two amplitudes: amplitude A, with  $|A| = 1$  and amplitude B with  $|B| = 1$  and phase difference  $\phi_W = 90^\circ$  between A and B that is entirely due to phase factors in CKM matrix elements.  
Draw the vector-addition diagram for the total amplitude A+B and calculate the magnitude  $|A + B|$
- (b) Now draw the diagram for the CP-conjugate process. (What happens to  $\phi_W$  under CP conjugation?)
- (c) Calculate the magnitude of  $|A + B|$ . Is it different from  $|\overline{A + B}|$ ? Is there observable CP violation in this process?
- (d) Redo the exercise with the following modification: the phase difference between A and B is now  $\phi = \phi_s + \phi_w$ , where  $\phi_w = 90^\circ$  is the phase difference due to CKM factors and  $\phi_s = 45^\circ$ , which is due to other other physics in amplitudes A and B that is invariant under CP conjugation (typically final state interactions from the strong interaction).