



**SIGHTSEEING  
IN THE  
LANDSCAPE**

# CONTENTS

- Landscape remarks  
(physics/06041340, Dutch version 1998)
- RCFT orientifolds  
(with Huiszoon, Fuchs, Schweigert, Walcher)
- 2003-2004 results  
(with Dijkstra, Huiszoon)
- 2005-2006 results  
(with Anastasopoulos, Dijkstra, Kiritsis, hep-th/0605226)

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- Beginning of last century:  
Einstein + Maxwell theory.  
Suggest a unique underlying unified theory.

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- Beginning of last century:  
Einstein + Maxwell theory.  
Suggest a unique underlying unified theory.
- Then some experimental problems arise:
  - Strong and Weak interactions
  - Muon (quark/lepton families)
  - Parameters (masses, couplings)

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- Beginning of last century:  
Einstein + Maxwell theory.  
Suggest a unique underlying unified theory.
- Then some experimental problems arise:
  - Strong and Weak interactions
  - Muon (quark/lepton families)
  - Parameters (masses, couplings)
- Then some theoretical problems arise:  
Yang-Mills theory: QED is not unique.  
Many other gauge theories are possible.

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- The Standard Model is discovered  
Once again suggests an underlying unified theory.  
(gauge principle; GUT structure). But uniqueness??

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- The Standard Model is discovered  
Once again suggests an underlying unified theory.  
(gauge principle; GUT structure). But uniqueness??
- String Theory is discovered.  
Unifies all interactions with gravity.  
Imposes strong restrictions on matter:  
Renewed hopes for uniqueness.

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- The Standard Model is discovered  
Once again suggests an underlying unified theory.  
(gauge principle; GUT structure). But uniqueness??
- String Theory is discovered.  
Unifies all interactions with gravity.  
Imposes strong restrictions on matter:  
Renewed hopes for uniqueness.
- The Duality Revolution of 1995:  
String Theory (M-Theory) is unique.  
(if we can define it...)

# UNIFICATION / UNIQUENESS

## A BRIEF HISTORY

- The Standard Model is discovered  
Once again suggests an underlying unified theory.  
(gauge principle; GUT structure). But uniqueness??
- String Theory is discovered.  
Unifies all interactions with gravity.  
Imposes strong restrictions on matter:  
Renewed hopes for uniqueness.
- The Duality Revolution of 1995:  
String Theory (M-Theory) is unique.  
(if we can define it...)
- But there is another revolution most people preferred to overlook: The string vacuum revolution.

**1984-2006:  
A SLOW REVOLUTION**

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions

A. Strominger,  
“Calabi-Yau manifolds with Torsion”, 1986

**All predictive power seems to have been lost.**

**All of this points to the overwhelming need to find a dynamical principle for determining the ground state, which now appears more imperative than ever.**

Lerche, Lüst, Schellekens

“Chiral, Four-dimensional Heterotic Strings From Self-Dual Lattices”, 1986

$(\Gamma_{22} \times D_3 \times (D_7)^9)_L$ , a Euclidean lattice of dimension 88. A lower limit on the total number of such lattices is provided by the Siegel mass formula [21] [22]

**this number is of order  $10^{1500}$  !**

It seems that not much is left of the once celebrated uniqueness of string theory.

Even if all that string theory could achieve would be a completely finite theory of all interactions including gravity, but with no further restrictions on the gauge groups and the representations, it would be a considerable success. But the situation

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions
- 1987: Gepner models
- .....
- 1995: M-theory compactifications, F-theory, Orientifolds
- .....

M.Dine

hep-th/0402101


Faced with this plethora of states, I, for a long time, comforted myself that not a single example of a (meta)stable ground state of this sort had been exhibited in a controlled approximation, and so perhaps there might be some unique or at least limited set of sensible states.

# A CERN CAFETARIA NAPKIN (~ 1988)



*All gauge theories*


# A CERN CAFETARIA NAPKIN (~ 1988)



● Complexity

All gauge theories


# A CERN CAFETARIA NAPKIN (~ 1988)



- Complexity
- Life

All gauge theories

# A CERN CAFETARIA NAPKIN (~ 1988)





- Complexity
- Life
- Intelligence

All gauge theories

# A CERN CAFETARIA NAPKIN (~ 1988)


Unique vacuum



- Complexity
- Life
- Intelligence

All gauge theories


# A CERN CAFETARIA NAPKIN (~ 1988)



- Complexity
- Life
- Intelligence

All gauge theories

# A CERN CAFETARIA NAPKIN (~ 1988)



- Complexity
- Life
- Intelligence

All gauge theories

# Naar een waardig slot

*Bert Schellekens*

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions
- 1987: Gepner models
- .....
- 1995: M-theory compactifications, F-theory, Orientifolds
- .....

# 1984-2006: A SLOW REVOLUTION

- 1984: Hopes for Unification and Uniqueness
- 1985: Calabi-Yau manifolds, Orbifolds, Narain Lattices.
- 1986: CY's with torsion; Fermionic and Bosonic constructions
- 1987: Gepner models
- .....
- 1995: M-theory compactifications, F-theory, Orientifolds
- .....
- 2003: "The Anthropic Landscape of String Theory" (L. Susskind)

# THE ANTHROPIIC PRINCIPLE

# THE ANTHROPIIC PRINCIPLE

- Most formulations are nonsense. (including statements by Brandon Carter, Barrows & Tipler).

# THE ANTHROPIIC PRINCIPLE

- Most formulations are nonsense. (including statements by Brandon Carter, Barrows & Tipler).
- Does not make sense without String Theory (or better) or Eternal Inflation (or equivalent).

# THE ANTHROPIIC PRINCIPLE

- Most formulations are nonsense. (including statements by Brandon Carter, Barrows & Tipler).
- Does not make sense without String Theory (or better) or Eternal Inflation (or equivalent).
- Is an inevitable consequence of String Theory.

# THE ANTHROPIIC PRINCIPLE

- Most formulations are nonsense. (including statements by Brandon Carter, Barrows & Tipler).
- Does not make sense without String Theory (or better) or Eternal Inflation (or equivalent).
- Is an inevitable consequence of String Theory.
- Until 2000, almost no papers relate String Theory and the Anthropic principle.

# THE ANTHROPIIC PRINCIPLE

- Most formulations are nonsense. (including statements by Brandon Carter, Barrows & Tipler).
- Does not make sense without String Theory (or better) or Eternal Inflation (or equivalent).
- Is an inevitable consequence of String Theory.
- Until 2000, almost no papers relate String Theory and the Anthropic principle.
- Without anti-anthropocentric prejudices, we might have predicted the “Anthropic Landscape of Quantum Gravity”.

# HINDSIGHT...

Soon after starting graduate school, I went to see Howard Georgi. “What are you thinking about?” he asked me. I rattled off several things that seemed interesting to me, ending with, “... and quantum gravity.” “**Don’t waste your time!**” he barked, “There’s no decoupling limit in which it’s sensible to consider quantum gravity effects, while neglecting other interactions. Unless you know particle physics all the way up to the Planck scale, you can never hope to say anything predictive about quantum gravity.” Howard was, of course, completely correct.

*Jacques Distler, “Musings”*

# HOW MANY “VACUA” ARE NEEDED?

- Requires understanding of “anthropic” considerations for different gauge theories.
- Requires some definition of a measure and boundaries.

Wild guess: about  $10^{20}$  for SM fine-tunings

The same problems exist in principle for the cosmological constant, but seem less serious there: about  $10^{120}$  would be needed.

Recent estimates: String Theory has plenty of ground states to understand all fine-tunings.

*(Bousso-Polchinski, Douglas Denef,...)*

# VACUUM COUNTING (1998)

$$10^{30} \times 10^{-80} = 10^{-50}$$

Number of vacua

SM Probability  
(experimental)

# VACUUM COUNTING (2006)

$$10^{500} \times 10^{-80} \times 10^{-120} = 10^{300}$$

↑  
Number of vacua

↑  
SM Probability

↑  
Cosmological  
Constant

# FALSIFIABILITY

# FALSIFIABILITY

- String Theory is falsifiable (but, remarkably, has not been falsified yet).

# FALSIFIABILITY

- String Theory is falsifiable (but, remarkably, has not been falsified yet).
  - Chiral Fermions (without anomalies)
  - The Standard Model gauge group
  - Three Families
  - Couplings of reasonable size
  - Two loop finiteness
  - Black hole entropy
  - Cosmological constant
  - Moduli stabilization
  - ....

# FALSIFIABILITY

- String Theory is falsifiable (but, remarkably, has not been falsified yet).
  - Chiral Fermions (without anomalies)
  - The Standard Model gauge group
  - Three Families
  - Couplings of reasonable size
  - Two loop finiteness
  - Black hole entropy
  - Cosmological constant
  - Moduli stabilization
  - ....
- Its vacuum structure is (theoretically) falsifiable.

# FALSIFIABILITY

- String Theory is falsifiable (but, remarkably, has not been falsified yet).
  - Chiral Fermions (without anomalies)
  - The Standard Model gauge group
  - Three Families
  - Couplings of reasonable size
  - Two loop finiteness
  - Black hole entropy
  - Cosmological constant
  - Moduli stabilization
  - ....
- Its vacuum structure is (theoretically) falsifiable.
- Non-anthropoc nature of other vacua is (theoretically) falsifiable.

# SUMMARY:

# SUMMARY:

- A landscape of vacua is the only sensible outcome for a “Theory of Everything”

# SUMMARY:

- A landscape of vacua is the only sensible outcome for a “Theory of Everything”
- Therefore: A Success for String Theory

# SUMMARY:

- A landscape of vacua is the only sensible outcome for a “Theory of Everything”
- Therefore: A Success for String Theory
- 4-D Quantum gravity implies that the SM is part of a huge landscape: an amazing conclusion! (if correct).

# SUMMARY:

- A landscape of vacua is the only sensible outcome for a “Theory of Everything”
- Therefore: A Success for String Theory
- 4-D Quantum gravity implies that the SM is part of a huge landscape: an amazing conclusion! (if correct).
- Fits nicely with some of the great discoveries in the history of science (heliocentric model, theory of Evolution...)



# Demystification by huge numbers:

- Planets (Giordano Bruno)
- Mutations (Evolution)
- Universes (Eternal Inflation)
- Alternative “Standard Models” (The Landscape)

## Demystification by huge numbers:

- Planets (Giordano Bruno)
- Mutations (Evolution)
- Universes (Eternal Inflation)
- Alternative “Standard Models” (The Landscape)

## A repetition of an old mistake:

There is nothing “special” about us.

This line of thought fits in very well with a series of insights that pointed out our modest place in the cosmos. Our planet is not the center of the solar system, our sun is just one of many stars and not even a very special one, and the same is true for our galaxy. It seems natural to assume that also our universe, including the quarks, leptons and interactions we observe is just one out of many possibilities.

(From physics/06041340)



● String Theory has never looked better...

● String Theory has never looked better...

● ... but it has never looked harder.

# REASONABLE GOALS

# REASONABLE GOALS

- Explore unknown regions of the landscape

# REASONABLE GOALS

- Explore unknown regions of the landscape
- Establish the likelihood of standard model features  
(gauge group, three families, ....)

# REASONABLE GOALS

- Explore unknown regions of the landscape
- Establish the likelihood of standard model features (gauge group, three families, ....)
- Convince ourselves that the standard model is a plausible vacuum.

# REASONABLE GOALS

- Explore unknown regions of the landscape
- Establish the likelihood of standard model features (gauge group, three families, ....)
- Convince ourselves that the standard model is a plausible vacuum.
- Determine if we are the “Chinese” or the “Andorrans” of the landscape.

# REASONABLE GOALS

- Explore unknown regions of the landscape
- Establish the likelihood of standard model features (gauge group, three families, ....)
- Convince ourselves that the standard model is a plausible vacuum.
- Determine if we are the “Chinese” or the “Andorrans” of the landscape.
- ... and maybe we get lucky



ORIENTIFOLDS  
OF  
GEPNER MODELS

# EARLIER FOOTPRINTS

C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y. S. Stanev, Phys. Lett. B **387** (1996) 743 [arXiv:hep-th/9607229].

R. Blumenhagen and A. Wisskirchen, Phys. Lett. B **438**, 52 (1998) [arXiv:hep-th/9806131].

G. Aldazabal, E. C. Andres, M. Leston and C. Nunez, JHEP **0309**, 067 (2003) [arXiv:hep-th/0307183].

I. Brunner, K. Hori, K. Hosomichi and J. Walcher, arXiv:hep-th/0401137.


R. Blumenhagen and T. Weigand, JHEP **0402** (2004) 041 [arXiv:hep-th/0401148].

G. Aldazabal, E. C. Andres and J. E. Juknevich, JHEP **0405**, 054 (2004) [arXiv:hep-th/0403262].

# THE LONG ROAD TO THE CHIRAL SSM

- ✿ Angelantonj, Bianchi, Pradisi, Sagnotti, Stanev (1996)  
*Chiral spectra from Orbifold-Orientifolds*
- ✿ Aldazabal, Franco, Ibanez, Rabadan, Uranga (2000)  
Blumenhagen, Görlich, Körs, Lüst (2000)  
Ibanez, Marchesano, Rabadan (2001)  
*Non-supersymmetric SM-Spectra with RR tadpole cancellation*
- ✿ Cvetič, Shiu, Uranga (2001)  
*Supersymmetric SM-Spectra with chiral exotics*
- ✿ Blumenhagen, Görlich, Ott (2002)  
Honecker (2003)  
*Supersymmetric Pati-Salam Spectra with brane recombination*
- ✿ Dijkstra, Huiszoon, Schellekens (2004)  
*Supersymmetric Standard Model (Gepner Orientifolds)*
- ✿ Honecker, Ott (2004)  
*Supersymmetric Standard Model ( $Z_6$  orbifold/orientifold)*

# CLOSED STRING PARTITION FUNCTION




$$P(\tau, \bar{\tau}) = \sum_{ij} \chi_i(\tau) Z_{ij} \chi_j(\bar{\tau})$$

Type IIB


# ORIENTIFOLD PARTITION FUNCTIONS

# ORIENTIFOLD PARTITION FUNCTIONS

$\frac{1}{2}$




+




# ORIENTIFOLD PARTITION FUNCTIONS


$\frac{1}{2}$




+





$\frac{1}{2}$




+



# ORIENTIFOLD PARTITION FUNCTIONS



# TRANSVERSE CHANNEL



# GEPNER MODELS

Building Blocks:  
Minimal N=2 CFT

$$c = \frac{3k}{k+2}, \quad k = 1, \dots, \infty$$

168 ways of solving  $\sum_i c_{k_i} = 9$

Spectrum:

$$h_{l,m} = \frac{l(l+2) - m^2}{4(k+2)} + \frac{s^2}{8}$$

$$(l = 0, \dots, k; \quad q = -k, \dots, k+2; \quad s = -1, 0, 1, 2)$$

(plus field identification)

$4(k+2)$  simple currents

# TENSORING

- ✱ Preserve world-sheet susy
- ✱ Preserve space-time susy (GSO)
- ✱ Use surviving simple currents to build MIPFs
- ✱ This yields one point in the moduli space of a Calabi-Yau manifold

# SELECTING MIPFs AND ORIENTIFOLDS

Each tensor product has a discrete group  $\mathcal{G}$   
of simple currents:  $J \cdot a = b$

Choose:

- ✱ A subgroup  $\mathcal{H}$  of  $\mathcal{G}$
- ✱ A rational matrix  $X_{\alpha\beta}$  defined on  $\mathcal{H}$

- ✱ An element  $K$  of  $\mathcal{G}$
- ✱ A set of signs  $\beta_K(J)$  defined on  $\mathcal{H}$

# A MIPF

$$\begin{aligned} & (0+2)^2 + (1+3)^2 + (4+6)*(13+15) + (5+7)*(12+14) \\ & + (8+10)^2 + (9+11)^2 + (12+14)*(5+7) + (13+15)*(4+6) \\ & + (16+18)*(25+27) + (17+19)*(24+26) + (20+22)^2 + (21+23)^2 \\ & + (24+26)*(17+19) + (25+27)*(16+18) + (28+30)^2 + (29+31)^2 \\ & + (32+34)^2 + (33+35)^2 + (36+38)*(45+47) + (37+39)*(44+46) \\ & + (40+42)^2 + (41+43)^2 + (44+46)*(37+39) + (45+47)*(36+38) \\ & + (48+50)*(57+59) + (49+51)*(56+58) + (52+54)^2 + (53+55)^2 \\ & + (56+58)*(49+51) + (57+59)*(48+50) + (60+62)^2 + (61+63)^2 \end{aligned}$$

....

$$\begin{aligned} & + 2*(2913)*(2915) + 2*(2914)*(2912) + 2*(2915)*(2913) \\ & + 2*(2916)^2 + 2*(2917)^2 + 2*(2918)^2 + 2*(2919)^2 \\ & + 2*(2920)^2 + 2*(2921)^2 + 2*(2922)^2 + 2*(2923)^2 \\ & + 2*(2924)*(2926) + 2*(2925)*(2927) + 2*(2926)*(2924) \\ & + 2*(2927)*(2925) + 2*(2928)^2 + 2*(2929)^2 + 2*(2930)^2 \\ & + 2*(2931)^2 + 2*(2932)*(2934) + 2*(2933)*(2935) \\ & + 2*(2934)*(2932) + 2*(2935)*(2933) + 2*(2936)*(2938) \\ & + 2*(2937)*(2939) + 2*(2938)*(2936) + 2*(2939)*(2937) \\ & + 2*(2940)^2 + 2*(2941)^2 + 2*(2942)^2 + 2*(2943)^2 \end{aligned}$$

# BOUNDARIES AND CROSSCAPS\*

## ☀ Boundary coefficients

$$R_{[a, \psi_a](m, J)} = \sqrt{\frac{|\mathcal{H}|}{|\mathcal{C}_a| |\mathcal{S}_a|}} \psi_a^*(J) S_{am}^J$$

## ☀ Crosscap coefficients

$$U_{(m, J)} = \frac{1}{\sqrt{|\mathcal{H}|}} \sum_{L \in \mathcal{H}} e^{\pi i (h_K - h_{KL})} \beta_K(L) P_{LK, m} \delta_{J, 0}$$

\*Huiszoon, Fuchs, Schellekens, Schweigert, Walcher (2000)

# COEFFICIENTS

## ☼ Klein bottle

$$K^i = \sum_{m, J, J'} \frac{S_m^i U_{(m, J)} g_{J, J'}^{\Omega, m} U_{(m, J')}}{S_{0m}}$$

## ☼ Annulus

$$A_{[a, \psi_a][b, \psi_b]}^i = \sum_{m, J, J'} \frac{S_m^i R_{[a, \psi_a]}(m, J) g_{J, J'}^{\Omega, m} R_{[b, \psi_b]}(m, J')}{S_{0m}}$$

## ☼ Moebius

$$M_{[a, \psi_a]}^i = \sum_{m, J, J'} \frac{P_m^i R_{[a, \psi_a]}(m, J) g_{J, J'}^{\Omega, m} U_{(m, J')}}{S_{0m}}$$

$$g_{J, J'}^{\Omega, m} = \frac{S_{m0}}{S_{mK}} \beta_K(J) \delta_{J', J^c}$$

# PARTITION FUNCTIONS

## ☀ Closed

$$\frac{1}{2} \left[ \sum_{ij} \chi_i(\tau) Z_{ij} \chi_i(\bar{\tau}) + \sum_i K_i \chi_i(2\tau) \right]$$

## ☀ Open

$$\frac{1}{2} \left[ \sum_{i,a,n} N_a N_b A^i_{ab} \chi_i\left(\frac{\tau}{2}\right) + \sum_{i,a} N_a M^i_a \hat{\chi}_i\left(\frac{\tau}{2} + \frac{1}{2}\right) \right]$$

$N_a$ : Chan-Paton multiplicity

# TADPOLES & ANOMALIES

- ✱ Tadpole cancellation condition:

$$\sum_b N_b R_{b(m,J)} = 4\eta_m U_{(m,J)}$$

- ✱ Cubic  $\text{Tr}F^3$  anomalies cancel


- ✱ Remaining anomalies by Green-Schwarz mechanism

- ✱ In rare cases, additional conditions for global anomaly cancellation\*


\*Gato-Rivera, Schellekens (2005)

# ABELIAN MASSES

Green-Schwarz mechanism



Axion-Vector boson vertex



Generates mass vector bosons of anomalous symmetries

(*e.g.*  $B + L$ )

But may also generate mass for non-anomalous ones

( $Y, B - L$ )

# SCOPE OF THE SEARCH

# SCOPE OF THE SEARCH

✻ 168 Gepner models

# SCOPE OF THE SEARCH

☼ 168 Gepner models

☼ 5403 MIPFs

# SCOPE OF THE SEARCH

☀ 168 Gepner models

☀ 5403 MIPFs

☀ 49322 Orientifolds

# SCOPE OF THE SEARCH

- ✻ 168 Gepner models
- ✻ 5403 MIPFs
- ✻ 49322 Orientifolds
- ✻  $45761187347637742772$  combinations of four boundary labels (brane stacks)


# SCOPE OF THE SEARCH

- ✻ 168 Gepner models
- ✻ 5403 MIPFs
- ✻ 49322 Orientifolds
- ✻ 45761187347637742772 combinations of four boundary labels (brane stacks)

Essential to decide what to search for!

# WHAT TO SEARCH FOR

## The Madrid model



Chiral  $SU(3) \times SU(2) \times U(1)$  spectrum:


$$3(u, d)_L + 3u_L^c + 3d_L^c + 3(e^-, \nu)_L + 3e_L^+$$

Y massless  $Y = \frac{1}{6}Q_a - \frac{1}{2}Q_c - \frac{1}{2}Q_d$

N=1 Supersymmetry

No tadpoles, global anomalies

# THE HIDDEN SECTOR



# REQUIRED SPECTRUM

3 families of  $SU(3) \times SU(2) \times U(1)$

+ non-chiral matter

# STATISTICS

Total number of 4-stack configurations	45761187347637742772 ( $45.7 \times 10^{18}$ )
Total number scanned	43752168618082181524
Total number of SM configurations	45051902 fraction: $1.0 \times 10^{-12}$
Total number of tadpole solutions	1649642 fraction: $3.8 \times 10^{-14}$ (*)
Total number of distinct solutions	211634

(\*) cf. Gmeiner, Blumenhagen, Honecker, Lüst, Weigand: "One in a Billion"

Standard model type: 6  
 Number of factors in hidden gauge group: 0  
 Gauge group: U(3) x Sp(2) x U(1) x U(1)

Number of representations: 19


3 x (V ,V ,0 ,0 ) chirality 3  
 3 x (V ,0 ,V ,0 ) chirality -3  
 3 x (V ,0 ,V\*,0 ) chirality -3  
 9 x (0 ,V ,0 ,V ) chirality 3  
 5 x (0 ,0 ,V ,V ) chirality -3  
 3 x (0 ,0 ,V ,V\*) chirality -3  
 2 x (V ,0 ,0 ,V )  
 10 x (0 ,V ,V ,0 )  
 2 x (Ad,0 ,0 ,0 )  
 2 x (A ,0 ,0 ,0 )


.....

Higgs:	(2,1/2)+ 2*,1/2)				5
Non-chiral SM matter	(Q,U,D,L,E,N):	0	0	0	3 1 0
Adjoint:		2	0	9	3
Symmetric Tensors:		1	10	7	3
Anti-Symmetric Tensors:		1	14	3	2
Lepto-quarks:	3,-1/3), 3,2/3)			1	0
Non-SM	a,b,c,d)	0	0	0	0
Hidden	Total dimension)	0			(chirality 0)

$$\sin^2(\theta_w) = .5271853$$


$$\frac{\alpha_3}{\alpha_2} = 3.2320501$$





# SU(5) MODELS

U(5)



# SU(5)

Type:	U	0	0	
Dimension	5	1	1	
3 x	(A	,0	,0	) chirality 3
11 x	(V	,V	,0	) chirality -3
8 x	(S	,0	,0	) chirality 0
3 x	(Ad,	0	,0	) chirality 0
1 x	(0	,A	,0	) chirality 0
3 x	(0	,V	,V	) chirality 0
8 x	(V	,0	,V	) chirality 0
2 x	(0	,S	,0	) chirality 0
4 x	(0	,0	,S	) chirality 0
4 x	(0	,0	,A	) chirality 0

*Note: gauge group is just SU(5)!*

# SUMMARY

Examples exist of chiral orientifold SSM spectra exist

- Without mirrors
- Without adjoints
- Without (anti)-symmetric tensors
- Without Observable-Hidden matter
- Without hidden sector

# SUMMARY

Examples exist of chiral orientifold SSM spectra exist

- Without mirrors
- Without adjoints
- Without (anti)-symmetric tensors
- Without Observable-Hidden matter
- Without hidden sector

...but to get all this simultaneously requires more statistics



IT'S JUST ONE SMALL STEP:  
874 HODGE NUMBERS SCANNED  
AT LEAST 30000 KNOWN (M. KREUZER)